

COORDINATED SCIENCE LABORATORY

AD690128

**EFFECT OF A BOUNDARY
ON THE BEAM
PLASMA INSTABILITY**

KENNETH EVANS, JR.
E. ATLEE JACKSON

DDT
REC
JUL 23 1969
A

UNIVERSITY OF ILLINOIS - URBANA, ILLINOIS

**EFFECT OF A BOUNDARY ON THE
BEAM PLASMA INSTABILITY**

Kenneth Evans, Jr. and E. Atlee Jackson

**Department of Physics and Coordinated Science Laboratory
University of Illinois, Urbana, Illinois**

**This work was supported in whole by Joint Services Electronics
Program (U.S. Army, U.S. Navy, and U.S. Air Force) under Contract DAAB
07-67-C-0199.**

**Reproduction in whole or in part is permitted for any purpose of
the United States Government.**

**This document has been approved for public release and sale; its
distribution is unlimited.**

**EFFECT OF A BOUNDARY ON THE
BEAM PLASMA INSTABILITY***

Kenneth Evans, Jr. and E. Atlee Jackson

Department of Physics and Coordinated Science Laboratory

University of Illinois, Urbana, Illinois

In recent years there has been considerable experimental¹⁻⁵ and some theoretical^{6,7} interest in the spatial-temporal growth of instabilities arising when a beam is injected into a plasma. There does not appear, however, to be any theoretical treatment of the modification of the growth of an initial perturbation due to the presence of a spatial boundary. This modification will be illustrated for the case of a monoenergetic electron beam injected into a cold plasma with an infinitely massive ion background.

The treatment is based on the linearized hydrodynamic equations and Maxwell's equations for the normalized densities, the drift velocities, and the electric field, ($\mathcal{E} = (e/m) E$):

$$n_{P,t} + u_{P,x} = 0 \quad (1)$$

$$u_{P,t} + \mathcal{E} = -\nu u_P \quad (2)$$

$$n_{B,t} + u_o n_{B,x} + u_{B,x} = 0 \quad (3)$$

$$u_{B,t} + u_o u_{B,x} + \mathcal{E} = 0 \quad (4)$$

$$\mathcal{E}_{,t} - \omega_P^2 u_P - \omega_B^2 (u_B + u_o n_B) = 0 \quad (5)$$

Eq. (2) contains a phenomenological collision term, representing collisions between the plasma electrons and the ion background. The remaining Maxwell's equation:

$$\mathcal{E}_{,x} + \omega_P^2 n_P + \omega_B^2 n_B = 0 \quad (6)$$

is treated as a condition to be satisfied at $t = 0$, whence it is satisfied for all later times.

Eqs. (1-5), which are of the following form (repeated indices are summed):

$$a_{ij}f_{j,t} + b_{ij}f_{j,x} + c_{ij}f_j = 0 \quad i, j = 1, \dots, 5 \quad (7)$$

are to be solved for $x > 0$ and $t > 0$ subject to conditions: $f_i(0, t) = g_i(t)$ and $f_i(x, 0) = h_i(x)$. These conditions are not all independent since Eqs. (2) and (5) must be satisfied at $x = 0$, and equation (6) must be satisfied at $t = 0$. The first two of these restrictions are due to the fact that characteristics of Eqs. (1-5) lie on the boundary $x = 0$.

The double Laplace transformation solution to Eqs. (1-5) is:

$$f_i(x, t) = (2\pi i)^{-2} \int dp dq e^{pt + qx} F_i(p, q) \quad (8)$$

$$F_i(p, q) = [a_{jk}H_k(q) + b_{jk}G_k(p)] d_{ji}(p, q)/D(p, q) \quad (9)$$

H_i and G_i are the one dimensional Laplace transforms of $h_i(x)$ and $g_i(t)$. D is the determinant of the matrix $a_{ij}p + b_{ij}q + c_{ij}$, and d_{ij} is the cofactor. ($D = 0$ is the dispersion relation, apart from a multiplicative factor p , for the infinite case.) The integration paths in (8) are to the right of all singularities of the integrand.

If the transforms H_k contain only simple poles at $q = \alpha_i$, $i = 1, \dots$, the q integration in Eq. (8) may be performed to give:

$$f_i(x, t) = \frac{1}{2\pi i} \sum_{l=1,2} \int dp e^{pt + q_l x} \frac{[a_{jk}H_k(q_l) + b_{jk}G_k(p)] d_{ji}(p, q_l)}{(\partial D(p, q)/\partial q)_{q=q_l}} + \frac{1}{2\pi i} \sum_m \int dp e^{pt + \alpha_m x} \frac{a_{jk}d_{ji}(p, \alpha_m)}{D(p, \alpha_m)} \text{Residue}(H_k(\alpha_m)) \quad (10)$$

where $q_l(p)$ are the roots of $D(p, q(p)) = 0$:

$$q_{1,2}(p) = \frac{-p}{u_0} \pm \frac{i\omega_B}{u_0} \frac{p(p + v)}{p(p + v) + \omega_p^2}^{\frac{1}{2}}$$

If $x > u_0 t$, the first term in Eq. (10) does not contribute, since the contour may be closed at infinity in the right half-plane, where it encloses no singularities. (This result is independent of whether the H_k have only simple poles.) The expression for $f_1(x, t)$ is then independent of the presence of the boundary and consists, as in the unbounded case, of plane wave terms proportional to $\exp(p_\ell(\alpha_1)t + \alpha_1 x)$ where $p_\ell(q)$, $\ell = 1, \dots, 4$ are the roots of $D(p(q), q) = 0$.

If $x < u_0 t$, the first term of Eq. (10) has singularities at: (a) singularities of $q_\ell(p)$, (b) singularities of $H_k(q_\ell(p))$ apart from those in (a), and (c) singularities of $G_k(p)$, which we assume to be simple poles at $p = \beta_i$, $i = 1, \dots$. The contour can, in this case, be closed in the left half-plane, and it can be shown that the singularities (b) exactly cancel the second term in Eq. (10). This cancellation eliminates the plane wave terms which appear for the infinite plasma. In their place the singularities (a), which include the essential singularities at $p = -\frac{1}{2}v \pm i[\omega_p^2 - \frac{1}{2}v^2]^{\frac{1}{2}}$, give the more complicated dependence due to the presence of the boundary. The singularities (c) give rise to plane wave terms proportional to $\exp(\beta_i t + q_\ell(\beta_i)x)$.

To illustrate the long time behavior of an initial perturbation, consider the case: $n_p(x, 0) = N_0 \sin kx$ and $\mathcal{E}(x, 0) = -N_0 \omega_p^2 k^{-1} (1 - \cos kx)$. All other boundary and initial conditions are taken to be zero. The integral (10) around the essential singularity can be evaluated by the saddle point method in the limit of large t . It can be shown that this method is valid in the present case, and that only two of the eight saddle points which occur

contribute to the asymptotic behavior. The result is:

$$\begin{aligned} \mathcal{E}(x,t) = & N_0 k u_0^2 (6\pi)^{-1/2} (\omega_p/\omega_B)^2 \beta^{1/2} \chi^{1/3} \tau^{-5/6} \\ & \times \exp \left[\frac{3}{4} \sqrt{3} \beta \chi^{2/3} \tau^{1/3} - (\frac{1}{2} v/\omega_p) \tau \right] \\ & \times \sin \left[\beta^{-3} \tau - \frac{3}{4} \beta \chi^{2/3} \tau^{1/3} - \phi - \frac{1}{12} \pi \right] \end{aligned} \quad (12)$$

where $\beta = [1 - (\frac{1}{2} v/\omega_p)^2]^{-1/6}$, $\chi = \omega_B x/u_0$, $\tau = \omega_p t$, and $\sin \phi = \frac{1}{2} v/\omega_p$. This result is valid provided $x \ll u_0 t$ and $\beta \chi^{2/3} \tau^{-2/3} \ll 1 \ll \beta \chi^{2/3} \tau^{1/3}$. It should be noted that the exponential factor, which dominates the form of $\mathcal{E}(x,t)$, is independent of the initial disturbance.

The above results yield the following picture for the development of an initial disturbance. If the disturbance is unstable according to the usual theory for spatially unbounded systems, it grows as $e^{\gamma t}$ only until $t = x/u_0$. For $t \gg x/u_0$ the amplitude goes as in Eq. (12). It reaches a maximum at $t_0 = \omega_p^{-1/2} (\frac{1}{2} \sqrt{3} \beta/v)^{3/2} (\omega_B x/u_0)$ and finally decays as $e^{-\frac{1}{2} v t}$. If the collision frequency is large, t_0 does not occur in the asymptotic region $t \gg x/u_0$, so the behavior is always $e^{-\frac{1}{2} v t}$ in this region. On the other hand if $v = 0$, the disturbance grows indefinitely, and one has essentially an absolute rather than a convective instability. This is due to the fact that there are characteristics of Eqs. (1-5) which are parallel to the t axis.⁸ The presence of collisions thereby insures the validity of the linear approximation for all times, provided x is not too large.

In experimental measurements what is observed is the result of many disturbances originating at different prior times. Since each disturbance has a limited lifetime (for any fixed x and $v \neq 0$), the cumulative

effect is a stationary state which varies with x . In the present situation the lifetime depends critically on the collision frequency, so one expects the level of fluctuations in the stationary state to also be strongly influenced by ν . While a rough estimate of this stationary state can be obtained from the above results, a more accurate analysis is desirable. Such an analysis is presently being performed.

One of the authors (A.J.) would like to express his sincere appreciation for the hospitality and financial assistance of the FOM-Instituut voor Plasma-Fysica, Rijnhuizen, Jutphaas, Netherlands.

*This work was supported in part by the Joint Services Electronics Program (U.S. Army, U.S. Navy, and U.S. Air Force) under contract DAAB-07-67-C-0199.

¹S.M. Levitskii and I.P. Shashurin, JETP 52, 350 (1967), Soviet Phys. JETP 25, 227 (1967).

²H.J. Hopman and W. Ott, Plasma Phys. 10, 315 (1968).

³H. Böhmer and M. Raether, IXth International Conference on Phenomena in Ionized Gases (Bucharest, 1969) and Proceeding of the Conference on Non-linear Effects in Plasmas (University of Paris, Orsay, 1966) to be published in Kalman and Feix, Non-linear Effects in Plasmas (Gordon and Breach, London)

⁴J.R. Apel, Phys. Fluids 12, 291, 640 (1969).

⁵S.A. Self, Stanford University Institute for Plasma Research Reports 275 and 276 (1969).

⁶W.E. Drummond, Phys. Fluids 7, 816 (1964).

⁷Ya. B. Fainberg and V.D. Shapiro, JETP 47, 1389 (1964), Soviet Phys. JETP 20, 937 (1965).

⁸A.I. Akhiezer, et. al., Collective Oscillations in a Plasma (The M.I.T. Press, Cambridge, 1967), Chap. II, Sec. 13.

Security Classification

DOCUMENT CONTROL DATA - R & D

(Security classification of title, body or abstract and indexing annotation must be entered when the overall report is classified)

1. ORIGINATING ACTIVITY (Corporate author) University of Illinois Coordinated Science Laboratory Urbana, Illinois 61801		2a. REPORT SECURITY CLASSIFICATION Unclassified	
		2b. GROUP	
3. REPORT TITLE EFFECT OF A BOUNDARY ON THE BEAM PLASMA INSTABILITY			
4. DESCRIPTIVE NOTES (Type of report and inclusive dates)			
5. AUTHOR(S) (First name, middle initial, last name) EVANS, Kenneth, Jr, & JACKSON, E. Atlee			
6. REPORT DATE July, 1969		7a. TOTAL NO. OF PAGES 6	7b. NO. OF REFS 8
8a. CONTRACT OR GRANT NO. DAAB 07-67-C-0199		9a. ORIGINATOR'S REPORT NUMBER(S) R-422	
b. PROJECT NO.		9b. OTHER REPORT NO(S) (Any other numbers that may be assigned this report)	
c.			
d.			
10. DISTRIBUTION STATEMENT This document has been approved for public release and sale; its distribution is unlimited.			
11. SUPPLEMENTARY NOTES		12. SPONSORING MILITARY ACTIVITY Joint Services Electronics Program thru U.S. Army Electronics Command Fort Monmouth, New Jersey 07703	
13. ABSTRACT NONE			

DD FORM 1 NOV 65 1473

Security Classification

Security Classification

14 KEY WORDS	LINK A		LINK B		LINK C	
	ROLE	WT	ROLE	WT	ROLE	WT
Plasma-beam instability Boundary effects						

Security Classification